



State of Utah  
Department of Human  
Services  
Division of Substance  
Abuse and Mental Health

# Student Health And Risk Prevention

## 2013 Prevention Needs Assessment Survey Revised Results

Report Prepared By:  
Bach Harrison, L.L.C.  
116 South 500 East  
Salt Lake City,  
Utah 84102  
Phone: 801-359-2064

**Weber & Morgan Counties  
LSAA  
DSAMH Region Profile Report  
(Revised January 2014)**

# Table of Contents

---

- Introduction
- The Charts and Tables in this Report
- Data Charts:
  - *Lifetime and 30 Day ATOD Use*
  - *Problem Substance Use, Mental Health, and Antisocial Behavior*
  - *Sources of Alcohol and Places of Alcohol Use*
  - *Risk and Protective Factor Profiles*
- The Risk and Protective Factor Model of Prevention
- Building a Strategic Prevention Framework
- School and Community Improvement Using Survey Data
- Risk and Protective Factor Scale Definitions
- Data Tables
- Appendix
- Contacts for Prevention

## Revision Note:

*SHARP Survey reports were revised on 1-17-2014 to correct lifetime use rates for substances other than alcohol and tobacco. While the corrections mainly affect the results for grade 10, grade 12, and all grades combined, some very small changes may be seen in the rates for grades 6 and 8. These revised reports also contain lifetime and 30-day e-cigarette use rates, which were not included in previous reports. Should you have any questions on these changes or require technical assistance, please contact SHARP Project Director Mary Johnstun at [mary@bach-harrison.com](mailto:mary@bach-harrison.com) or 801-842-2682.*

# Introduction

## 2013 Weber & Morgan Counties LSAA Prevention Needs Assessment Survey Report

This report summarizes the findings from the Utah 2013 Prevention Needs Assessment (PNA) Survey that was conducted as part of the Student Health and Risk Prevention (SHARP) Statewide Survey. The survey was administered to students in grades 6, 8, 10 and 12 in 39 school districts and 14 charter schools across Utah.

The results for your Local Substance Abuse Authority (LSAA) region are presented along with comparisons to 2009 and 2011 SHARP Survey results, where applicable. Results from administrations prior to 2009 may be found by consulting past years' profile reports. The PNA Survey was designed to assess adolescent substance use, anti-social behavior, and the risk and protective factors that predict adolescent problem behaviors.

Further, in keeping with the vision that prevention services are designed to have a positive impact on the lives of individuals, efforts have been made to ensure that the PNA survey also gathers data on issues such as mental health and suicide, gang involvement, academic issues, health and fitness, and other

prevention-related topics.

Table 1 contains the characteristics of the students who completed the survey from your LSAA and the State of Utah. Because not all students answer all of the questions, the total number of survey respondents by gender and survey respondents by ethnicity may be less than the reported total students.

When using the information in this report, please pay attention to the number of students who participated from your community. If **60% or more** of the students participated, the report is a good indicator of the levels of substance use, risk, protection, and antisocial behavior. If fewer than 60% participated, consult with your local prevention coordinator or a survey professional before generalizing the results to the entire community.

Coordination and administration of the Utah PNA Survey was a collaborative effort of State of Utah, Department of Human Services, Division of Substance Abuse and Mental Health; Office of Education; Department of Health; and Bach Harrison, LLC. For more information about the PNA or prevention services in Utah, please refer to the *Contacts for Prevention* section at the end of this report.

**Table 1. Characteristics of Participants**

Total Survey Respondents	LSAA 2009		LSAA 2011		LSAA 2013		State 2013	
	Number	Percent	Number	Percent	Number	Percent	Number	Percent
	3,160	100	4,350	100	3,763	100	47,137	100
<b>Survey Respondents by Grade</b>								
6	968	30.6	1,349	31.0	1,153	30.6	13,923	29.5
8	945	29.9	1,300	29.9	1,089	28.9	14,040	29.8
10	774	24.5	957	22.0	791	21.0	10,816	22.9
12	473	15.0	744	17.1	730	19.4	8,358	17.7
<b>Survey Respondents by Gender</b>								
Male	1,529	49.1	2,160	49.8	1,873	49.9	22,760	48.4
Female	1,585	50.9	2,177	50.2	1,877	50.1	24,218	51.6
<b>Survey Respondents by Ethnicity</b>								
African American	52	1.7	77	1.8	90	2.4	742	1.6
Asian	48	1.5	53	1.2	45	1.2	750	1.6
Hispanic	531	17.0	711	16.5	661	17.7	6,029	13.0
American Indian	52	1.7	93	2.2	71	1.9	838	1.8
Pacific Islander	28	0.9	38	0.9	37	1.0	697	1.5
White	2,198	70.3	2,985	69.4	2,449	65.6	33,612	72.4
Multi-racial	219	7.0	345	8.0	381	10.2	3,731	8.0

# The Charts and Tables in this Report

There are five types of charts presented in this report: 1) substance use charts, 2) problem use, mental health and antisocial behavior (ASB) charts, 3) sources of alcohol acquisition, 4) places of alcohol consumption, 5) risk factor charts and 6) protective factor charts. Data from the charts are presented numerically in Tables 3 through 9.

## Understanding the Format of the Charts

There are several graphical elements common to all the charts. Understanding the format of the charts and what these elements represent is essential in interpreting the results of the 2013 SHARP survey.

- **The Bars** on substance use and antisocial behavior charts represent the percentage of students in that grade who reported a given behavior. The bars on the risk and protective factor charts represent the percentage of students whose answers reflect significant risk or protection in that category.

Each set of differently colored bars represents one of the last three administrations of the PNA: 2009, 2011, and 2013. By looking at the percentages over time, it is possible to identify trends in substance use and antisocial behavior. By studying the percentage of youth at risk and with protection over time, it is possible to determine whether the percentage of students at risk or with protection is increasing, decreasing, or staying the same. This information is important when deciding which risk and protective factors warrant attention.

- **Dots and Diamonds** provide points of comparison to larger samples. The dots on the charts represent the percentage of all of the youth surveyed across Utah who reported substance use, problem behavior, elevated risk, or elevated protection.

For the 2013 PNA Survey, there were 47,137 participants in grades 6, 8, 10, and 12, out of 72,709 sampled, a participation rate of 64.8%. The fact that over 47,000 students across the state participated in the PNA make the state dot a good estimate of the rates of ATOD use and levels of risk and protective factors of youth in Utah. The survey results provide considerable information for communities to use in planning prevention services.

The diamonds represent national data from either the Monitoring the Future (MTF) survey or the Bach Harrison Norm. The Bach Harrison Norm was developed by Bach Harrison LLC to provide states

and communities with the ability to compare their results on risk, protection, and antisocial measures with more national measures. Survey participants from eight statewide surveys and five large regional surveys across the nation were combined into a database of approximately 460,000 students. The results were weighted to make the contribution of each state and region proportional to its share of the national population. Bach Harrison analysts then calculated rates for antisocial behavior and for students at risk and with protection. The results appear on the charts as BH Norm. In order to keep the Bach Harrison Norm relevant, it is updated approximately every two years as new data become available.

A comparison to state-wide and national results provides additional information for your community in determining the relative importance of levels of alcohol, tobacco and other drug (ATOD) use, antisocial behavior, risk, and protection. Information about other students in the state and the nation can be helpful in determining the seriousness of a given level of problem behavior. Scanning across the charts, it is important to observe the factors that differ the most from the Bach Harrison Norm. This is the first step in identifying the levels of risk and protection that are higher or lower than those in other communities. The risk factors that are higher than the Bach Harrison Norm and the protective factors that are lower than the Bach Harrison Norm are factors your community should consider addressing when planning prevention programs.

## Lifetime and 30-Day ATOD Use

- **Lifetime use** is a measure of the percentage of students who tried the particular substance at some point in their lifetime and is used to show the percentage of students who have had experience with a particular substance.
- **30-day use** is a measure of the percentage of students who used the substance at least once in the 30 days prior to taking the survey and is a more sensitive indicator of the level of current use of the substance.

## Problem Substance Use, Need for Treatment, and Antisocial Behavior

- **Problem substance use** is measured in several different ways: binge drinking (having five or more drinks in a row during the **two weeks prior**

## The Charts and Tables in this Report

to the survey), use of one-half a pack or more of cigarettes **per day**, and youth indicating drinking alcohol and driving or reporting riding with a driver who had been drinking alcohol during **the past 30 days**.

- **The need for treatment** estimates the percentage of students in need of substance abuse and mental health treatment.

**Substance abuse treatment needs** are estimates of youth in need of alcohol treatment, drug treatment and an estimate of students that need either alcohol OR drug treatment. The need for treatment is defined as students who have used alcohol on 10 or more occasions in their lifetime or any drugs in their lifetime and marked three or more of the following six items related to their **drug or alcohol use in the past year**: 1) spent more time using than intended, 2) neglected some of your usual responsibilities because of use, 3) wanted to cut down on use, 4) others objected to your use, 5) frequently thought about using, and 6) used alcohol or drugs to relieve feeling such as sadness, anger, or boredom. Students could mark each item as it related to their drug and/or alcohol use.

**Needs Mental Health Treatment** was estimated using the K6 Scale that was developed with support from the National Center for Health Statistics for use in the National Health Interview Survey. The tool screens for psychological distress by asking students “During **the past 30 days**, how often did you: 1) feel nervous? 2) feel hopeless? 3) feel restless or fidgety? 4) feel so depressed that nothing could cheer you up? 5) feel that everything was an effort? and 6) feel worthless?”

Answers were scored based on responses: *None of the time* (0 points), *A little of the time* (1 point), *Some of the time* (2 points), *Most of the time* (3 points), *All of the time* (4 points). Students with a score of 13 or more points were determined to be in need of mental health treatment.

**Youth considering suicide** are also in need of mental health services. This section of the report also contains the percentage of students answering yes to the question, “During **the past 12 months**, did you ever seriously consider attempting suicide?”

- **Antisocial behavior (ASB)** is a measure of the percentage of students who report any involvement **during the past year** with the eight antisocial behaviors listed in the charts.

### Sources of Alcohol and Places of Alcohol Use

These charts present the percentage of students who obtained alcohol from nine specific sources and the percentage that used alcohol in six specific places during the past year. Questions regarding sources of alcohol were not asked in 2011, but were included on the 2013 SHARP PNA. The number of students reporting use is presented to assist in interpreting the results.

### Risk and Protective Factors

Risk and protective factor scales measure specific aspects of a youth’s life experience that predict whether he/she will engage in problem behaviors. The scales, defined in Table 2, are grouped into four domains: community, family, school, and peer/individual. The risk and protective factor charts show the percentage of students at risk and with protection for each of the scales.

### Additional Tables in this Report

Tables 10 to 12 contain additional data for prevention planning and reporting to state and federal agencies.

### Drug Free Communities

Table 10 contains information relevant to Drug Free Community (DFC) grantees. These tables report the four DFC Core Measures on alcohol, tobacco, marijuana and prescription drugs:

- **Perception of Risk** - The percentage of respondents who report that regular use of the substance has *moderate risk* or *great risk*
- **Perception of Parental/Peer Disapproval** - The percentage of respondents who report their parents feel regular use of alcohol/ANY use of cigarettes, marijuana, or prescription drugs is *wrong* or *very wrong*.
- **Past 30-Day Use** - The percentage surveyed reporting using the substance at least once in the past 30 days

### Data for Prevention Planning


Table 11 contains information on student perceptions of school safety, bullying, classroom and school discipline, and student perception of ATOD use among their peers.

### Perceived Parental Approval and ATOD Use

Table 12 explores the relationship between perceived parental approval and ATOD use. A full explanation of how to interpret these data is available accompanying the tables.

# Substance Use and Antisocial Behavior


## Lifetime and 30-Day ATOD Use\* 2013 Weber & Morgan Counties LSAA Student Survey, Grade 6



\* 2013 SHARP PNA lifetime use is calculated differently than previous years. Lifetime use values for substances other than alcohol and tobacco were updated January 2014.  
 \*\* The "Sedatives" question from 2009 was split into "Prescription Tranquilizers" and "Prescription Sedatives" in 2011. "E-Cigarette" data (available for 2011 and 2013) have been added in the January 2014 revision.  
 † "Prescription Drugs" is a combined measure showing the total use of any "Stimulant," "Sedative," "Tranquilizer," or "Narcotic Prescription" drugs (Consult appendix for explanation of above changes.)  
 †† Monitoring the Future does not survey 6th grade students.

# Substance Use and Antisocial Behavior

## Lifetime and 30-Day ATOD Use\* 2013 Weber & Morgan Counties LSAA Student Survey, Grade 8




\* 2013 SHARP PNA lifetime use is calculated differently than previous years. Lifetime use values for substances other than alcohol and tobacco were updated January 2014.

\*\* The "Sedatives" question from 2009 was split into "Prescription Tranquilizers" and "Prescription Sedatives" in 2011. "E-Cigarette" data (available for 2011 and 2013) have been added in the January 2014 revision.

† "Prescription Drugs" is a combined measure showing the total use of any "Stimulant," "Sedative," "Tranquilizer," or "Narcotic Prescription" drugs (Consult appendix for explanation of above changes.)

# Substance Use and Antisocial Behavior


## Lifetime and 30-Day ATOD Use\* 2013 Weber & Morgan Counties LSAA Student Survey, Grade 10



\* 2013 SHARP PNA lifetime use is calculated differently than previous years. Lifetime use values for substances other than alcohol and tobacco were updated January 2014.  
 \*\* The "Sedatives" question from 2009 was split into "Prescription Tranquilizers" and "Prescription Sedatives" in 2011. "E-Cigarette" data (available for 2011 and 2013) have been added in the January 2014 revision.  
 † "Prescription Drugs" is a combined measure showing the total use of any "Stimulant," "Sedative," "Tranquilizer," or "Narcotic Prescription" drugs (Consult appendix for explanation of above changes.)

# Substance Use and Antisocial Behavior


## Lifetime and 30-Day ATOD Use\* 2013 Weber & Morgan Counties LSAA Student Survey, Grade 12



\* 2013 SHARP PNA lifetime use is calculated differently than previous years. Lifetime use values for substances other than alcohol and tobacco were updated January 2014.  
 \*\* The "Sedatives" question from 2009 was split into "Prescription Tranquilizers" and "Prescription Sedatives" in 2011. "E-Cigarette" data (available for 2011 and 2013) have been added in the January 2014 revision.  
 † "Prescription Drugs" is a combined measure of any "Stimulant," "Sedative," "Tranquilizer," or "Narcotic Prescription" drugs (Consult appendix for explanation of above changes.)

# Substance Use and Antisocial Behavior

## Lifetime and 30-Day ATOD Use\* 2013 Weber & Morgan Counties LSAA Student Survey, All Grades



\* 2013 SHARP PNA lifetime use is calculated differently than previous years. Lifetime use values for substances other than alcohol and tobacco were updated January 2014.


\*\* The "Sedatives" question from 2009 was split into "Prescription Tranquilizers" and "Prescription Sedatives" in 2011. "E-Cigarette" data (available for 2011 and 2013) have been added in the January 2014 revision.

† "Prescription Drugs" is a combined measure showing the total use of any "Stimulant," "Sedative," "Tranquilizer," or "Narcotic Prescription" drugs (Consult appendix for explanation of above changes.)

†† "All Grades" MTF data are not available.

# Substance Use and Antisocial Behavior

## Problem Substance Use, Need For Treatment, and Antisocial Behavior 2013 Weber & Morgan Counties LSAA Student Survey, Grade 6




\* 6th grade MTF data for "Binge Drinking" and "1/2 Pack Cigarettes/Day" are unavailable.

\*\* Due to a change in how lifetime drug use was measured, the calculation for "Needs Drug Treatment" changed slightly. (See appendix for details.)

† National comparison data for "Driving While Drinking," "Riding with a Drinking Driver," and "Antisocial Behavior" are Bach Harrison Norm values. Please see Tables 5 and 6 for more information on the time frames for the values presented in this chart.

# Substance Use and Antisocial Behavior

## Problem Substance Use, Need For Treatment, and Antisocial Behavior 2013 Weber & Morgan Counties LSAA Student Survey, Grade 8




\* National comparison data for "Binge Drinking" and "1/2 Pack Cigarettes/Day" are Monitoring the Future values.

\*\* Due to a change in how lifetime drug use was measured, the calculation for "Needs Drug Treatment" changed slightly. (See appendix for details.)

† National comparison data for "Driving While Drinking," "Riding with a Drinking Driver," and "Antisocial Behavior" are Bach Harrison Norm values. Please see Tables 5 and 6 for more information on the time frames for the values presented in this chart.

# Substance Use and Antisocial Behavior

## Problem Substance Use, Need For Treatment, and Antisocial Behavior 2013 Weber & Morgan Counties LSAA Student Survey, Grade 10




\* National comparison data for "Binge Drinking" and "1/2 Pack Cigarettes/Day" are Monitoring the Future values.

\*\* Due to a change in how lifetime drug use was measured, the calculation for "Needs Drug Treatment" changed slightly. (See appendix for details.)

† National comparison data for "Driving While Drinking," "Riding with a Drinking Driver," and "Antisocial Behavior" are Bach Harrison Norm values. Please see Tables 5 and 6 for more information on the time frames for the values presented in this chart.

# Substance Use and Antisocial Behavior

## Problem Substance Use, Need For Treatment, and Antisocial Behavior 2013 Weber & Morgan Counties LSAA Student Survey, Grade 12




\* National comparison data for "Binge Drinking" and "1/2 Pack Cigarettes/Day" are Monitoring the Future values.

\*\* Due to a change in how lifetime drug use was measured, the calculation for "Needs Drug Treatment" changed slightly. (See appendix for details.)

† National comparison data for "Driving While Drinking," "Riding with a Drinking Driver," and "Antisocial Behavior" are Bach Harrison Norm values. Please see Tables 5 and 6 for more information on the time frames for the values presented in this chart.

# Substance Use and Antisocial Behavior

## Problem Substance Use, Need For Treatment, and Antisocial Behavior 2013 Weber & Morgan Counties LSAA Student Survey, All Grades



\* "All Grades" MTF data for "Binge Drinking" and "1/2 Pack Cigarettes/Day" are unavailable.


\*\* Due to a change in how lifetime drug use was measured, the calculation for "Needs Drug Treatment" changed slightly. (See appendix for details.)

† National comparison data for "Driving While Drinking," "Riding with a Drinking Driver," and "Antisocial Behavior" are Bach Harrison Norm values. Please see Tables 5 and 6 for more information on the time frames for the values presented in this chart.

# Sources of Alcohol

## Sources Of Alcohol\*

2013 Weber & Morgan Counties LSAA Student Survey, Grade 6




\* Questions regarding sources of alcohol were not asked in 2011.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Sources of Alcohol

## Sources Of Alcohol\* 2013 Weber & Morgan Counties LSAA Student Survey, Grade 8




\* Questions regarding sources of alcohol were not asked in 2011.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Sources of Alcohol

## Sources Of Alcohol\* 2013 Weber & Morgan Counties LSAA Student Survey, Grade 10




\* Questions regarding sources of alcohol were not asked in 2011.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Sources of Alcohol

## Sources Of Alcohol\* 2013 Weber & Morgan Counties LSAA Student Survey, Grade 12




\* Questions regarding sources of alcohol were not asked in 2011.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Sources of Alcohol

## Sources Of Alcohol\* 2013 Weber & Morgan Counties LSAA Student Survey, All Grades




\* Questions regarding sources of alcohol were not asked in 2011.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Places of Alcohol Use

## Places Of Alcohol Use 2013 Weber & Morgan Counties LSAA Student Survey, Grade 6




\*"At or near school" is new for the 2011 SHARP PNA.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Places of Alcohol Use

## Places Of Alcohol Use 2013 Weber & Morgan Counties LSAA Student Survey, Grade 8




\*\*At or near school" is new for the 2011 SHARP PNA.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Places of Alcohol Use

## Places Of Alcohol Use 2013 Weber & Morgan Counties LSAA Student Survey, Grade 10




\*\*At or near school" is new for the 2011 SHARP PNA.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Places of Alcohol Use

## Places Of Alcohol Use 2013 Weber & Morgan Counties LSAA Student Survey, Grade 12




\*\*At or near school" is new for the 2011 SHARP PNA.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Places of Alcohol Use

## Places Of Alcohol Use 2013 Weber & Morgan Counties LSAA Student Survey, All Grades




\*\*At or near school" is new for the 2011 SHARP PNA.

\*\* Sample size represents the number of youth who chose at least one place of drinking alcohol. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

# Risk and Protective Factor Profiles

## Risk Profile


2013 Weber & Morgan Counties LSAA Student Survey, Grade 6



# Risk and Protective Factor Profiles

## Protective Profile


2013 Weber & Morgan Counties LSAA Student Survey, Grade 6



# Risk and Protective Factor Profiles

## Risk Profile


2013 Weber & Morgan Counties LSAA Student Survey, Grade 8



# Risk and Protective Factor Profiles

## Protective Profile


2013 Weber & Morgan Counties LSAA Student Survey, Grade 8



# Risk and Protective Factor Profiles

## Risk Profile


2013 Weber & Morgan Counties LSAA Student Survey, Grade 10



# Risk and Protective Factor Profiles

## Protective Profile


2013 Weber & Morgan Counties LSAA Student Survey, Grade 10



# Risk and Protective Factor Profiles

## Risk Profile


2013 Weber & Morgan Counties LSAA Student Survey, Grade 12



# Risk and Protective Factor Profiles

## Protective Profile


2013 Weber & Morgan Counties LSAA Student Survey, Grade 12



# Risk and Protective Factor Profiles

## Risk Profile


2013 Weber & Morgan Counties LSAA Student Survey, All Grades



# Risk and Protective Factor Profiles

## Protective Profile

2013 Weber & Morgan Counties LSAA Student Survey, All Grades



# The Risk and Protective Factor Model

Prevention is a science. The Risk and Protective Factor Model of Prevention is a proven way of reducing substance abuse and its related consequences. This model is based on the simple premise that to prevent a problem from happening, we need to identify the factors that increase the risk of that problem developing and then find ways to reduce the risks. Just as medical researchers have found risk factors for heart disease such as diets high in fat, lack of exercise, and smoking; a team of researchers at the University of Washington have defined a set of risk factors for youth problem behaviors.

**Risk factors** are characteristics of school, community and family environments, and of students and their peer groups known to predict increased likelihood of drug use, delinquency, school dropout, and violent behaviors among youth. For example, children who live in disorganized, crime-ridden neighborhoods are more likely to become involved in crime and drug use than children who live in safe neighborhoods.

The chart below shows the links between the 19 risk factors and five problem behaviors. The check marks indicate where at least two well designed, published research studies have shown a link between the risk factor and the problem behavior.

**Protective factors** exert a positive influence and buffer against the negative influence of risk, thus reducing the likelihood that adolescents will engage in problem behaviors. Protective factors identified through research include strong bonding to family, school, community and peers, and healthy beliefs and clear standards for behavior. Protective bonding depends on three conditions:

- **Opportunities** for young people to actively contribute
- **Skills** to be able to successfully contribute
- **Consistent recognition** or reinforcement for their efforts and accomplishments

Bonding confers a protective influence only when there is a positive climate in the bonded community. Peers and adults in these schools, families and neighborhoods must communicate healthy values and set clear standards for behavior in order to ensure a protective effect. For example, strong bonds to antisocial peers would not be likely to reinforce positive behavior.

Research on risk and protective factors has important implications for children’s academic success, positive youth development, and prevention of health and behavior problems. In order to promote academic success and positive youth development and to prevent problem behaviors, **it is necessary to address the factors that predict these outcomes.** By measuring risk and protective factors in a population, specific risk factors that are elevated and widespread can be identified and targeted by policies, programs, and actions shown to reduce those risk factors and to promote protective factors.

Each risk and protective factor can be linked to specific types of interventions that have been shown to be effective in either reducing risk(s) or enhancing protection(s). The steps outlined here will help your LSAA make key decisions regarding allocation of resources, how and when to address specific needs, and which strategies are most effective and known to produce results.

In addition to helping assess current conditions and prioritize areas of greatest need, data from the SHARP Prevention Needs Assessment (PNA) Survey can be a powerful tool in applying for and complying with several federal programs (such as the Strategic Prevention Framework process, the No Child Left Behind Act and Drug Free Communities grants), outlined later in this report. The survey also gathers valuable data which allows state and local agencies to address other prevention issues related to academic achievement, mental health, gang involvement, health and fitness, and personal safety.

Risk Factors for Adolescent Problem	Community							Family				School			Peer/Individual					
	Availability of Drugs	Availability of Firearms	Community Laws and Norms Favorable Toward Drug Use, Firearms and Crime	Media Portrayals of the Behavior	Transitions and Mobility	Low Neighborhood Attachment and Community Disorganization	Extreme Economic Deprivation	Family History of the Problem Behavior	Family Management Problems	Family Conflict	Favorable Parental Attitudes and Involvement in the Problem Behavior	Academic Failure Beginning in Late Elementary School	Lack of Commitment to School	Early and Persistent Antisocial Behavior	Rebelliousness	Gang Involvement	Friends Who Engage in the Problem Behavior	Favorable Attitudes Toward the Problem Behavior	Early Initiation of the Problem Behavior	Constitutional Factors
Substance Abuse	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Delinquency		✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Teen Pregnancy							✓	✓	✓	✓		✓	✓	✓			✓	✓	✓	
School Drop-Out					✓		✓	✓	✓			✓	✓	✓	✓		✓	✓	✓	
Violence	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Depression & Anxiety					✓			✓	✓	✓		✓		✓						✓

## Building a Strategic Prevention Framework

The PNA is an important data source for the Substance Abuse and Mental Health Services Administration (SAMHSA) Center for Substance Abuse Prevention (CSAP) Strategic Prevention Framework (SPF). CSAP created the SPF model to guide states and communities in creating planned, data-driven, effective, and sustainable prevention programs. Each part represents an interdependent element of the ongoing process of prevention coordination.

**Assessment:** Profile Population Needs, Resources, and Readiness to Address the Problems and Gaps in Service Delivery. The SPF begins with an assessment of the needs in the community that is based on data. The Utah State Epidemiological Outcomes Workgroup (SEOW) has compiled data from several sources to aid in the needs assessment process. One of the primary sources of needs assessment data is this Prevention Needs Assessment Survey (PNA). While planning prevention services, communities are urged to collect and use multiple data sources, including archival and social indicators, assessment of existing resources, key informant interviews, and community readiness. The PNA results presented in this Profile Report will help you to identify needs for prevention services. PNA data include adolescent substance use, anti-social behavior, and many of the risk and protective factors that predict adolescent problem behaviors.

**Capacity:** Mobilize and/or Build Capacity to Address Needs. Engagement of key stakeholders at the State and community levels is critical to plan and implement successful prevention activities that will be sustained over time. Some of the key tasks to mobilize the state and communities are to work with leaders and stakeholders to build coalitions, provide training, leverage resources, and help sustain prevention activities.

**Planning:** Develop a Comprehensive Strategic Plan.

States and communities should develop a strategic plan that articulates not only a vision for the prevention activities, but also strategies for organizing and implementing prevention efforts. The strategic plan should be based on the assessments conducted during Step 1. The Plan should address the priority needs, build on identified resources/strengths, set measurable objectives, and identify how progress will be monitored. Plans should be adjusted with ongoing needs assessment and monitoring activities.



# Building a Strategic Prevention Framework

**Implementation: Implement Evidence-based Prevention Programs and Infrastructure Development Activities.** By measuring and identifying the risk factors and other causal factors that contribute to the targeted problems specified in your strategic plan, programs can be implemented that will reduce the prioritized substance abuse problems. After completing Steps 1, 2, and 3, communities will be able to choose prevention strategies that have been shown to be effective, are appropriate for the population served, can be implemented with fidelity, are culturally appropriate, and can be sustained over time. One resource for evidence-based prevention practices is University Colorado at Boulder's Blueprints For Healthy Youth <http://www.colorado.edu/cspv/blueprints/>.

**Evaluation: Monitor Process, Evaluate Effectiveness, Sustain Effective Programs/Activities, and Improve or Replace Those That Fail:** Finally, ongoing monitoring and evaluation are essential to determine if the desired outcomes are achieved, assess service delivery quality, identify successes, encourage needed improvement, and promote sustainability of effective policies, programs, and practices. The PNA allows communities to monitor levels of ATOD use, antisocial behavior, risk, and protection.

**Sustainability and Cultural Competence** are at the core of the SPF model, indicating the key role they play in each of the five elements. Incorporating principles of cultural competence and sustainability throughout assessment, capacity appraisal, planning, implementation and evaluation helps ensure successful, long lasting prevention programs.

**Sustainability: Sustainability is accomplished by utilizing a comprehensive approach.** By building adaptive and flexible programs around a variety of resources, funding and organizations, states and communities will build sustainable programs and achieve sustainable outcomes. A strategic plan that dynamically responds to changing issues, data, priorities, and resources is more likely to achieve long term results.

Sharing information gathered during the evaluation stage with key stakeholders, forging partnerships and encouraging creative collaboration all enhance sustainability.

**Cultural Competence: Planners need to recognize the needs, styles, values and beliefs of the recipients of prevention efforts.** Culturally competent prevention strategies use interventions, evaluations and communication strategies appropriate to their intended community. Cultural issues reflect a range of influences and are not just a matter of ethnic or racial identity. Learning to communicate with audiences from diverse geographic, cultural, economic, social, and linguistic backgrounds can increase program efficacy and ensure sustainable results.

Whether enlisting extended family networks as a prevention resource for single parent households, or ensuring there are resources available to bridge language gaps, cultural competency will help you recognize differences in prevention needs and tailor prevention approaches accordingly.

A one-size-fits-all program is less effective than a program that draws on community-based values, traditions, and customs and works with knowledgeable people from the community to develop focused interventions, communication and support.

# School and Community Improvement Using Survey Data

## What are the numbers telling you?

- Review the charts and data tables presented in this report. Note your findings as you discuss the following questions.
- Which 3-5 risk factors appear to be higher than you would want when compared to the Bach Harrison Norm?
- Which 3-5 protective factors appear to be lower than you would want when compared to the Bach Harrison Norm?
- Which levels of 30-day drug use are increasing and/or unacceptably high?
  - Which substances are your students using the most?
  - At which grades do you see unacceptable usage levels?
- Which levels of antisocial behaviors are increasing and/or unacceptably high?
  - Which behaviors are your students exhibiting the most?
  - At which grades do you see unacceptable behavior levels?

## How to identify high priority problem areas

- **Look across the charts** – which items stand out as either much higher or much lower than the other?
- **Compare your data with statewide, and/or national data** – differences of 5% between local and other data are probably significant.
- **Prioritize problems for your area** – Make an assessment of the rates you’ve identified. Which can be realistically addressed with the funding available to your community? Which problems fit best with the prevention resources at hand?
- **Determine the standards and values held within your community** – For example: Is it acceptable in your community for a percentage of high school students to drink alcohol regularly as long as that percentage is lower than the overall state rate?

## Use these data for planning.

- **Substance use and antisocial behavior data** – raise awareness about the problems and promote dialogue.
- **Risk and protective factor data** – identify exactly where the community needs to take action.
- **Additional survey data** – use data on academic achievement, mental health and suicide, health and fitness, gang involvement, seat belt use, and other areas to broaden your prevention approach. Find ways to share these data with other prevention planners in your community.
- **Promising approaches** – access resources listed on the last page of this report for ideas about programs that have been proven effective in addressing the risk factors that are high in your community, and improving the protective factors that are low.

	Sample	Priority Rate 1	Priority Rate 2	Priority Rate 3
<b>Risk Factors</b>	6th grad Fav. Attitude to Drugs (Peer/Indiv. Scale) @ 15% (8% > BH Norm.)			
<b>Protective Factors</b>	10th grad - Rewards for prosocial involvm. (Schod Domain) (down 10% from 2 yrs ago)			
<b>30-day Substance Abuse</b>	8th grad Binge Drinking @ 13% (6% above state av.)			
<b>Antisocial Behavior</b>	12th grad - Drunk/High at School @ 21% (same as state, but still a priority.)			

## Risk and Protective Scale Definitions

**Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles**

<i>Community Domain Risk Factors</i>	
<i>Low Neighborhood Attachment</i>	Low neighborhood bonding is related to higher levels of juvenile crime and drug selling.
<i>Laws and Norms Favorable Toward Drug Use</i>	Research has shown that legal restrictions on alcohol and tobacco use, such as raising the legal drinking age, restricting smoking in public places, and increased taxation have been followed by decreases in consumption. Moreover, national surveys of high school seniors have shown that shifts in normative attitudes toward drug use have preceded changes in prevalence of use.
<i>Perceived Availability of Drugs and Handguns</i>	The availability of cigarettes, alcohol, marijuana, and other illegal drugs has been related to the use of these substances by adolescents. The availability of handguns is also related to a higher risk of crime and substance use by adolescents.
<i>Community Domain Protective Factors</i>	
<i>Rewards for Prosocial Involvement</i>	Rewards for positive participation in activities helps youth bond to the community, thus lowering their risk for substance use.
<i>Family Domain Risk Factors</i>	
<i>Poor Family Management</i>	Parents' use of inconsistent and/or unusually harsh or severe punishment with their children places them at higher risk for substance use and other problem behaviors. Also, parents' failure to provide clear expectations and to monitor their children's behavior makes it more likely that they will engage in drug abuse whether or not there are family drug problems.
<i>Family Conflict</i>	Children raised in families high in conflict, whether or not the child is directly involved in the conflict, appear at risk for both delinquency and drug use.
<i>Family History of Antisocial Behavior</i>	When children are raised in a family with a history of problem behaviors (e.g., violence or ATOD use), the children are more likely to engage in these behaviors.
<i>Parental Attitudes Favorable Toward Antisocial Behavior &amp; Drugs</i>	In families where parents use illegal drugs, are heavy users of alcohol, or are tolerant of children's use, children are more likely to become drug abusers during adolescence. The risk is further increased if parents involve children in their own drug (or alcohol) using behavior, for example, asking the child to light the parent's cigarette or get the parent a beer from the refrigerator.
<i>Family Domain Protective Factors</i>	
<i>Family Attachment</i>	Young people who feel that they are a valued part of their family are less likely to engage in substance use and other problem behaviors.
<i>Opportunities for Prosocial Involvement</i>	Young people who are exposed to more opportunities to participate meaningfully in the responsibilities and activities of the family are less likely to engage in drug use and other problem behaviors.
<i>Rewards for Prosocial Involvement</i>	When parents, siblings, and other family members praise, encourage, and attend to things done well by their child, children are less likely to engage in substance use and problem behaviors.
<i>School Domain Risk Factors</i>	
<i>Academic Failure</i>	Beginning in the late elementary grades (grades 4-6) academic failure increases the risk of both drug abuse and delinquency. It appears that the experience of failure itself, for whatever reasons, increases the risk of problem behaviors.
<i>Low Commitment to School</i>	Surveys of high school seniors have shown that the use of drugs is significantly lower among students who expect to attend college than among those who do not. Factors such as liking school, spending time on homework, and perceiving the coursework as relevant are also negatively related to drug use.
<i>School Domain Protective Factors</i>	
<i>Opportunities for Prosocial Involvement</i>	When young people are given more opportunities to participate meaningfully in important activities at school, they are less likely to engage in drug use and other problem behaviors.
<i>Rewards for Prosocial Involvement</i>	When young people are recognized and rewarded for their contributions at school, they are less likely to be involved in substance use and other problem behaviors.

## Risk and Protective Scale Definitions

**Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles (cont'd)**

<i>Peer-Individual Risk Factors</i>	
<i>Rebelliousness</i>	Young people who do not feel part of society, are not bound by rules, don't believe in trying to be successful or responsible, or who take an active rebellious stance toward society, are at higher risk of abusing drugs. In addition, high tolerance for deviance, a strong need for independence and normlessness have all been linked with drug use.
<i>Early Initiation of Antisocial Behavior and Drug Use</i>	Early onset of drug use predicts misuse of drugs. The earlier the onset of any drug use, the greater the involvement in other drug use and the greater frequency of use. Onset of drug use prior to the age of 15 is a consistent predictor of drug abuse, and a later age of onset of drug use has been shown to predict lower drug involvement and a greater probability of discontinuation of use.
<i>Attitudes Favorable Toward Antisocial Behavior and Drug Use</i>	During the elementary school years, most children express anti-drug, anti-crime, and pro-social attitudes and have difficulty imagining why people use drugs or engage in antisocial behaviors. However, in middle school, as more youth are exposed to others who use drugs and engage in antisocial behavior, their attitudes often shift toward greater acceptance of these behaviors. Youth who express positive attitudes toward drug use and antisocial behavior are more likely to engage in a variety of problem behaviors, including drug use.
<i>Perceived Risk of Drug Use</i>	Young people who do not perceive drug use to be risky are far more likely to engage in drug use.
<i>Interaction with Antisocial Peers</i>	Young people who associate with peers who engage in problem behaviors are at higher risk for engaging in antisocial behavior themselves.
<i>Friends' Use of Drugs</i>	Young people who associate with peers who engage in alcohol or substance abuse are much more likely to engage in the same behavior. Peer drug use has consistently been found to be among the strongest predictors of substance use among youth. Even when young people come from well-managed families and do not experience other risk factors, spending time with friends who use drugs greatly increases the risk of that problem developing.
<i>Rewards for Antisocial Behavior</i>	Young people who receive rewards for their antisocial behavior are at higher risk for engaging further in antisocial behavior and substance use.
<i>Depressive Symptoms</i>	Young people who are depressed are overrepresented in the criminal justice system and are more likely to use drugs. Survey research and other studies have shown a link between depression and other youth problem behaviors.
<i>Gang Involvement</i>	Youth who belong to gangs are more at risk for antisocial behavior and drug use.
<i>Peer-Individual Protective Factors</i>	
<i>Belief in the Moral Order</i>	Young people who have a belief in what is "right" or "wrong" are less likely to use drugs.
<i>Religiosity</i>	Young people who regularly attend religious services are less likely to engage in problem behaviors.
<i>Interaction with Prosocial Peers</i>	Young people who associate with peers who engage in prosocial behavior are more protected from engaging in antisocial behavior and substance use.
<i>Prosocial Involvement</i>	Participation in positive school and community activities helps provide protection for youth.
<i>Rewards for Prosocial Involvement</i>	Young people who are rewarded for working hard in school and the community are less likely to engage in problem behavior.

# Data Tables

**Table 3. Percentage of Students Who Used ATODs During Their Lifetime†**

How old were you when you first/Have you ever: (Students indicating any answer other than Never)		Grade 6				Grade 8				Grade 10				Grade 12				All Grades			
		LSAA 2009	LSAA 2011	LSAA 2013	State 2013																
Alcohol	had alcoholic beverages (beer, wine or hard liquor) to drink - more than just a few sips?	10.5	10.0	10.1	6.5	26.3	26.3	18.0	14.7	38.1	37.0	35.9	27.1	41.1	41.7	49.4	33.1	28.6	28.3	28.4	20.0
Cigarettes	smoked cigarettes?	5.2	4.8	5.8	4.0	13.5	13.7	11.8	9.3	22.4	22.5	23.5	15.9	21.4	26.8	28.4	19.2	15.3	16.7	17.5	11.9
E-Cigarettes*	used electronic cigarettes or e-cigarettes?	n/a	0.8	4.9	2.4	n/a	3.7	14.3	6.4	n/a	7.7	33.5	13.6	n/a	12.8	41.7	16.6	n/a	6.1	23.7	9.6
Chewing Tobacco	used smokeless tobacco (chew, snuff, plug, dipping tobacco, chewing tobacco)?	1.7	0.6	0.9	0.7	4.2	3.7	1.4	1.9	6.3	7.2	4.1	3.5	7.3	9.4	10.8	6.3	4.8	5.1	4.3	3.0
Marijuana	used marijuana (grass, pot) or hashish (hash, hash oil)?	1.7	0.8	2.6	1.6	9.9	8.9	10.1	8.5	19.6	20.5	27.3	19.3	19.0	26.9	35.2	24.4	12.2	13.9	18.8	13.2
Hallucinogens	used LSD (acid) or other hallucinogens (like PCP, mescaline, peyote, "shrooms" or psilocybin)?	0.5	0.2	0.3	0.2	2.3	2.2	2.4	1.3	3.9	5.1	4.1	3.3	4.1	9.1	7.7	5.6	2.7	4.0	3.7	2.6
Cocaine	used cocaine (like cocaine powder) or "crack" (cocaine in chunk or rock form)?	0.6	0.1	0.6	0.3	1.9	1.0	0.5	0.6	2.9	1.5	1.7	1.5	1.5	5.2	2.5	2.2	1.7	1.8	1.3	1.1
Inhalants	sniffed glue, breathed the contents of an aerosol spray can, or inhaled other gases or sprays, in order to get high?	6.8	5.7	6.9	4.4	10.2	9.5	7.7	5.9	7.2	8.2	6.9	5.7	8.5	9.0	6.5	5.2	8.2	8.1	7.0	5.3
Methamphetamines	used methamphetamines (meth, speed, crank, crystal meth)?	0.7	0.0	0.5	0.1	0.9	1.1	0.5	0.5	1.8	1.6	1.0	1.0	0.9	2.6	2.3	1.4	1.1	1.3	1.1	0.7
Prescription Stimulants	used prescription stimulants or amphetamines (such as Adderall, Ritalin, or Dexedrine) without a doctor telling you to take them?	0.6	0.9	0.5	0.6	1.3	3.5	1.4	1.5	3.1	5.5	4.2	4.0	3.7	7.0	9.2	5.9	2.2	4.1	3.8	2.9
Prescription Sedatives*	used prescription sedatives including barbiturates or sleeping pills (such as phenobarbital, Tuinal, Seconal, Ambien, Lunesta, or Sonata) without a doctor telling you to take them?	1.9	2.9	2.2	1.8	3.7	6.3	4.3	3.5	7.3	9.3	7.2	5.6	9.8	8.8	7.6	5.5	5.6	6.8	5.3	4.1
Prescription Tranquilizers	used prescription tranquilizers (such as Librium, Valium, Xanax, Ativan, Soma, or Klonopin) without a doctor telling you to take them?		0.0	0.7	0.3		0.5	1.0	0.6		1.8	3.0	1.9		1.0	5.9	3.1		0.8	2.7	1.5
Narcotic Prescription Drugs	used narcotic prescription drugs (such as OxyContin, methadone, morphine, codeine, Demerol, Vicodin, Percocet) without a doctor telling you to take them?	0.5	0.3	0.6	0.3	2.3	1.4	1.0	0.8	6.0	6.6	3.1	3.0	8.8	9.5	8.8	5.3	4.3	4.3	3.4	2.3
Any Prescription Drugs**	used prescription drugs (stimulants, sedatives, tranquilizers, or narcotics) without a doctor telling you to take them?	2.7	3.7	2.8	2.4	5.2	8.4	6.0	4.5	10.5	13.2	10.5	8.4	14.0	15.8	16.0	10.9	8.0	10.1	8.9	6.4
Heroin	used heroin?	0.3	0.2	0.1	0.1	1.0	1.0	0.5	0.5	1.4	1.0	1.0	0.7	0.9	1.2	1.0	0.8	0.9	0.8	0.6	0.5

\* In 2011, "Sedatives" was split into "Prescription Tranquilizers" and "Prescription Sedatives." 2011 and 2013 "E-Cigarette" data were added in January 2014.

\*\* "Any Prescription Drugs" is a combined measure showing the total use of any "Stimulant," "Sedative," "Tranquilizer," or "Narcotic Prescription" drugs.

† 2013 SHARP PNA lifetime use is calculated differently than previous years. Lifetime use values for substances other than alcohol and tobacco were updated January 2014. (See appendix for details.)

# Data Tables

**Table 4. Percentage of Students Who Used ATODs During the Past 30 Days†**

In the past 30 days, on how many occasions (if any) have you... (One or more occasions)		Grade 6				Grade 8				Grade 10				Grade 12				All Grades			
		LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013
Alcohol	had alcoholic beverages (beer, wine or hard liquor) to drink - more than just a few sips?	2.1	0.8	1.9	1.0	10.1	7.9	7.2	4.2	18.1	12.8	13.3	9.4	20.3	20.2	21.1	14.0	12.4	10.1	10.9	7.0
Cigarettes	smoked cigarettes?	0.8	0.2	0.9	0.5	3.8	3.3	3.2	1.8	7.1	7.6	4.9	3.9	5.1	9.3	8.7	5.0	4.1	5.0	4.5	2.7
E-Cigarettes*	used electronic cigarettes or e-cigarettes?	n/a	0.5	1.7	1.2	n/a	1.3	9.5	2.9	n/a	3.9	23.5	7.0	n/a	6.0	26.5	8.0	n/a	2.8	15.3	4.7
Chewing Tobacco	used smokeless tobacco (chew, snuff, plug, dipping tobacco, chewing tobacco)?	0.6	0.2	0.2	0.3	2.7	1.0	0.6	0.6	3.9	2.4	1.2	1.1	2.4	3.2	2.1	1.8	2.4	1.6	1.0	0.9
Marijuana	used marijuana (grass, pot) or hashish (hash, hash oil)?	0.7	0.3	1.1	0.6	5.0	4.1	5.3	4.2	10.3	10.0	12.1	9.1	7.7	12.6	14.4	9.9	5.8	6.6	8.2	5.8
Hallucinogens	used LSD (acid) or other hallucinogens (like PCP, mescaline, peyote, "shrooms" or psilocybin)?	0.4	0.1	0.0	0.0	0.6	1.1	0.9	0.5	1.7	1.8	0.8	0.9	1.2	3.0	0.9	1.3	0.9	1.4	0.7	0.7
Cocaine	used cocaine (like cocaine powder) or "crack" (cocaine in chunk or rock form)?	0.3	0.1	0.3	0.1	0.3	0.4	0.0	0.2	1.0	0.2	0.3	0.3	0.3	0.0	0.2	0.4	0.5	0.2	0.2	0.3
Inhalants	sniffed glue, breathed the contents of an aerosol spray can, or inhaled other gases or sprays, in order to get high?	2.5	1.1	2.8	2.0	2.5	3.2	3.2	2.1	1.3	1.2	1.1	1.2	1.2	0.6	0.4	0.5	1.9	1.6	1.9	1.5
Methamphetamines	used methamphetamines (meth, speed, crank, crystal meth)?	0.3	0.0	0.0	0.0	0.3	0.4	0.2	0.2	0.7	0.3	0.0	0.2	0.0	0.3	0.2	0.3	0.3	0.2	0.1	0.2
Prescription Stimulants	used prescription stimulants or amphetamines (such as Adderall, Ritalin, or Dexedrine) without a doctor telling you to take them?	0.3	0.1	0.2	0.1	0.5	1.2	0.3	0.4	0.6	2.1	1.0	1.2	1.2	2.4	1.5	1.3	0.6	1.4	0.8	0.7
Prescription Sedatives*	used prescription sedatives including barbiturates or sleeping pills (such as phenobarbital, Tuinal, Seconal, Ambien, Lunesta, or Sonata) without a doctor telling you to take them?	0.5	1.0	0.8	0.6	1.5	2.3	1.6	1.2	3.2	4.0	1.9	1.8	2.6	3.3	3.4	1.7	1.9	2.6	1.9	1.3
Prescription Tranquilizers	used prescription tranquilizers (such as Librium, Valium, Xanax, Ativan, Soma, or Klonopin) without a doctor telling you to take them?	0.0	0.0	0.3	0.0	0.5	0.5	0.4	0.2	1.7	0.9	0.6	0.6	1.0	1.8	0.8	0.8	0.8	0.8	0.8	0.4
Narcotic Prescription Drugs	used narcotic prescription drugs (such as OxyContin, methadone, morphine, codeine, Demerol, Vicodin, Percocet) without a doctor telling you to take them?	0.0	0.1	0.1	0.1	0.5	0.4	0.2	0.3	2.6	2.5	0.6	1.1	2.7	1.4	3.0	1.5	1.4	1.1	1.0	0.7
Any Prescription Drugs**	used prescription drugs (stimulants, sedatives, tranquilizers, or narcotics) without a doctor telling you to take them?	0.7	1.1	1.1	0.7	2.0	3.2	2.3	1.7	4.6	6.1	3.3	3.3	4.8	5.7	6.9	3.5	3.0	4.0	3.4	2.3
Heroin	used heroin?	0.1	0.1	0.0	0.0	0.7	0.5	0.3	0.2	0.5	0.0	0.0	0.2	0.3	0.2	0.1	0.2	0.4	0.2	0.1	0.2
Ecstasy	used MDMA ('X', 'E', or ecstasy)?	0.3	0.0	0.0	0.0	0.7	1.1	0.4	0.4	2.8	1.8	0.5	0.7	2.2	3.0	0.7	0.6	1.5	1.4	0.4	0.4
Steroids	used steroids or anabolic steroids (such as Anadrol, Oxandrin, Durabolin, Equipoise or Depotesterone)?	0.7	0.1	0.1	0.2	0.4	0.3	0.2	0.3	0.9	0.5	0.3	0.3	1.5	0.7	0.4	0.4	0.9	0.4	0.2	0.3

\* In 2011, "Sedatives" was split into "Prescription Tranquilizers" and "Prescription Sedatives." 2011 and 2013 "E-Cigarette" data were appended in January 2014.

\*\* "Any Prescription Drugs" is a combined measure showing the total use of any "Stimulant," "Sedative," "Tranquilizer," or "Narcotic Prescription" drugs.

† Since not all students answer all questions, the percentage of students reporting use in the past 30 days may be greater than the percentage reporting age of first use. (In 2013, age of first use became the new basis for calculating lifetime use for substances other than alcohol and tobacco. See appendix for details.)

# Data Tables

**Table 5. Percentage of Students With Problem Substance Use and Treatment Needs**

		Grade 6				Grade 8				Grade 10				Grade 12				All Grades			
		LSAA 2009	LSAA 2011	LSAA 2013	State 2013																
<b>Problem Use</b>																					
Binge Drinking*	How many times have you had 5 or more alcoholic drinks in a row in the past 2 weeks? (One or more times)	3.1	1.3	2.8	1.4	7.6	6.0	5.3	3.4	11.0	9.8	8.8	6.1	15.7	12.1	13.3	9.1	9.2	7.1	7.6	4.9
1/2 Pack of Cigarettes/Day	During the past 30 days, how many cigarettes did you smoke per day? (11 to 20 cigarettes, More than 20 cigarettes)	0.0	0.0	0.2	0.1	0.3	0.3	0.3	0.1	0.8	0.4	0.2	0.4	0.8	1.6	1.0	0.8	0.5	0.6	0.4	0.3
<b>Alcohol and Driving</b>																					
Drinking and Driving	During the past 30 days, how many times did you DRIVE a car or other vehicle when you had been drinking alcohol?	1.1	0.6	1.1	1.0	1.7	1.7	1.7	1.8	1.9	1.8	2.8	2.1	5.0	4.4	4.5	3.6	2.4	2.1	2.6	2.1
Riding with a Drinking Driver	During the past 30 days, how many times did you RIDE in a car or other vehicle driven by someone who had been drinking alcohol?	13.8	7.6	8.1	6.7	16.7	8.7	9.6	7.8	13.8	10.9	11.0	9.3	12.2	13.9	9.6	8.8	14.2	10.4	9.6	8.2
<b>Need for Treatment</b>																					
Needs Alcohol Treatment	Answered "Yes" to at least 3 alcohol treatment questions and has used alcohol on 10 or more occasions	0.2	0.2	0.2	0.1	1.6	2.1	2.1	1.0	6.3	4.4	3.7	3.1	6.0	8.4	6.2	4.2	3.6	3.9	3.1	2.1
Needs Drug Treatment**	Answered "Yes" to at least 3 drug treatment questions and has any lifetime drug use.	0.5	0.1	0.8	0.6	1.2	2.5	4.7	3.1	4.5	7.0	7.3	6.1	2.9	7.4	8.3	6.2	2.3	4.5	5.3	3.9
Alcohol or Drug Treatment	Needs alcohol, drug or alcohol AND drug treatment as per criteria above	0.6	0.3	1.0	0.7	2.1	3.9	5.5	3.5	8.9	8.7	8.9	7.3	7.3	12.2	10.9	8.1	4.8	6.6	6.6	4.8
Needs Mental Health Treatment	Scored 13 or more points on the K6 screening scale for psychological distress. (See text for further explanation.)	15.4	11.0	10.5	9.5	15.2	12.3	14.4	13.6	14.4	12.5	16.3	15.6	7.8	12.5	13.2	13.2	13.0	12.1	13.6	13.0
Considering Suicide	Answered "Yes" to "During the past 12 months, did you ever seriously consider attempting suicide?"	n/a	5.1	7.3	7.2	n/a	12.1	14.7	13.5	n/a	11.9	17.3	15.6	n/a	8.6	12.3	12.8	n/a	9.6	12.9	12.3

\* Since not all students answer all questions, the percentage of students reporting binge drinking may be greater than the percentage reporting 30-day alcohol use.

\*\* Due to a change in how lifetime drug use was measured, the calculation for "Needs Drug Treatment" changed slightly. (See appendix for details.)

# Data Tables

**Table 6. Percentage of Students With Antisocial Behavior**

How many times in the past year (12 months) have you: (One or more times)	Grade 6				Grade 8				Grade 10				Grade 12				All Grades			
	LSAA 2009	LSAA 2011	LSAA 2013	State 2013																
Been Drunk or High at School	2.6	1.7	2.3	1.3	8.0	6.0	5.1	4.3	13.0	13.1	13.3	9.8	13.9	14.1	14.6	10.7	9.2	8.6	8.8	6.4
Been Suspended from School	11.4	10.5	9.6	5.6	17.7	12.4	13.5	8.9	16.2	14.2	11.1	8.6	8.1	11.9	8.4	6.3	13.4	12.3	10.7	7.3
Sold Illegal Drugs	0.6	0.2	0.5	0.3	3.4	1.9	2.1	1.7	5.3	5.0	6.0	4.4	3.7	7.1	6.2	4.8	3.2	3.5	3.7	2.8
Stolen or Tried to Steal a Motor Vehicle	1.5	1.0	0.7	0.6	3.4	1.2	1.3	1.2	3.7	2.3	3.2	1.8	1.2	1.6	0.6	1.1	2.4	1.5	1.5	1.2
Been Arrested	1.9	0.9	1.7	1.1	5.6	2.9	3.0	2.8	5.6	4.6	3.4	3.3	4.1	4.0	3.0	3.5	4.3	3.1	2.8	2.6
Attacked Someone with the Idea of Seriously Hurting Them	9.7	9.7	7.8	6.1	12.3	10.9	8.8	7.3	10.5	8.5	6.2	6.2	8.0	5.7	3.4	4.8	10.1	8.8	6.6	6.1
Carried a Handgun	3.7	6.3	6.3	6.9	4.6	4.7	7.9	8.0	7.0	4.5	8.5	7.6	4.6	5.1	5.1	7.4	4.9	5.1	6.9	7.5
Carried a Handgun to School	0.8	0.4	0.5	0.3	0.6	0.6	0.1	0.3	1.0	0.5	0.5	0.4	0.3	0.2	0.0	0.6	0.7	0.4	0.3	0.4

# Data Tables

**Table 7. Sources and Places of Alcohol Use**

If you drank alcohol (not just a sip or taste) in the past year, how did you get it?*	Grade 6				Grade 8				Grade 10				Grade 12				All Grades			
	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013
<b>Sample size**</b>	88	n/a	74	603	235	n/a	142	1,642	284	n/a	234	2,381	190	n/a	302	2,338	797	n/a	752	6,964
I bought it myself from a store	6.8	n/a	4.1	2.8	4.7	n/a	4.9	3.6	4.9	n/a	3.8	3.7	12.1	n/a	6.3	7.8	6.8	n/a	5.1	5.0
I got it at a party	40.9	n/a	45.9	37.5	61.7	n/a	45.8	48.8	74.3	n/a	62.4	60.8	82.6	n/a	74.2	71.6	68.9	n/a	62.4	59.6
I gave someone else money to buy it for me	14.8	n/a	8.1	7.5	20.9	n/a	23.9	18.5	34.2	n/a	29.9	30.0	55.3	n/a	47.7	46.1	33.1	n/a	33.8	30.7
I got it from someone I know age 21 or older	37.5	n/a	35.1	27.5	49.4	n/a	49.3	40.4	58.5	n/a	52.1	50.9	73.7	n/a	65.2	64.1	57.1	n/a	55.2	50.8
I got it from someone I know under age 21	19.3	n/a	18.9	18.1	38.7	n/a	26.8	31.7	53.5	n/a	39.3	38.3	55.3	n/a	40.1	38.2	45.8	n/a	35.2	35.0
I got it from a family member or relative other than my parents	44.3	n/a	33.8	32.7	46.4	n/a	27.5	34.9	34.5	n/a	35.5	32.0	38.4	n/a	26.8	30.3	40.0	n/a	30.3	32.2
I got it from home with my parents' permission	31.8	n/a	29.7	33.2	25.5	n/a	32.4	28.9	26.4	n/a	28.2	27.6	26.8	n/a	25.5	25.9	26.9	n/a	28.1	27.8
I got it from home without my parents' permission	28.4	n/a	17.6	17.7	38.3	n/a	31.0	36.6	38.0	n/a	28.2	32.2	34.7	n/a	29.8	26.0	36.3	n/a	28.3	29.9
I got it another way	34.1	n/a	24.3	24.9	28.1	n/a	28.2	23.1	27.5	n/a	21.4	20.9	18.4	n/a	17.2	18.0	26.2	n/a	21.3	20.8
During the past year did you drink alcohol at any of the following places?	Grade 6				Grade 8				Grade 10				Grade 12				All Grades			
	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013	LSAA 2009	LSAA 2011	LSAA 2013	State 2013
<b>Sample size**</b>	89	106	89	790	234	303	161	1,889	268	288	256	2,559	185	275	311	2,428	776	972	817	7,666
At my home or someone else's home without any parent permission	39.3	33.0	32.6	36.3	59.8	53.1	53.4	56.4	64.2	67.0	68.4	64.7	71.9	68.4	72.7	67.8	61.9	59.4	63.2	60.7
At my home with my parent's permission	39.3	59.4	53.9	54.3	32.9	41.9	46.6	43.9	32.5	33.3	39.5	37.6	32.4	32.0	38.3	36.5	33.4	38.5	42.0	40.5
At someone else's home with their parent's permission	14.6	31.1	28.1	25.1	25.6	30.0	30.4	24.9	36.2	42.7	32.8	34.5	44.9	44.4	43.7	43.0	32.6	38.0	36.0	33.8
At or near school†	n/a	24.5	12.4	15.6	n/a	14.5	14.3	16.8	n/a	18.1	13.7	15.3	n/a	9.5	10.9	13.8	n/a	15.2	12.6	15.2
In a car	14.6	23.6	13.5	16.8	23.1	15.5	18.0	18.2	34.3	23.3	20.3	24.8	35.1	21.5	20.6	27.8	28.9	20.4	19.2	23.3
In some other place	43.8	34.0	39.3	35.9	34.6	38.9	40.4	36.1	29.9	34.4	38.3	36.0	33.5	37.1	33.8	38.5	33.8	36.5	37.1	36.8

\* Questions regarding sources of alcohol were not asked in 2011.

\*\* Sample size represents the number of youth who chose at least one place of alcohol consumption. Students who indicated they had not drunk alcohol in the past year are not included in the sample. In the case of smaller sample sizes, caution should be exercised before generalizing results to the entire community.

† "At or near school" was introduced on the 2011 SHARP PNA.

# Data Tables

**Table 8. Percentage of Students Reporting Risk**

Risk Factor	Grade 6				Grade 8				Grade 10				Grade 12				All Grades			
	LSAA 2009	LSAA 2011	LSAA 2013	State 2013																
<b>Community Domain</b>																				
Low Neighborhood Attachment	45.3	43.1	36.7	33.0	34.9	28.0	30.1	27.0	36.4	33.9	34.2	34.9	37.8	37.7	44.0	37.8	38.7	35.5	36.4	33.1
Laws & Norms Favor Drug Use	32.3	25.4	27.9	21.7	31.2	22.3	24.0	17.8	27.0	21.1	21.6	15.3	28.7	22.0	23.6	19.6	29.9	22.6	24.2	18.6
Perceived Availability of Drugs	33.2	33.8	28.7	26.0	34.6	28.8	28.7	24.6	39.3	35.1	35.4	26.6	38.4	33.4	32.7	29.4	36.2	32.7	31.5	26.6
Perceived Availability of Handguns	19.5	20.3	23.7	20.4	36.3	35.0	33.7	33.3	22.7	22.4	25.0	26.3	25.3	34.0	31.3	31.6	25.9	28.0	28.4	27.9
<b>Family Domain</b>																				
Poor Family Management	43.2	44.7	41.3	36.4	37.7	33.0	32.9	28.8	33.4	29.0	37.4	27.0	34.7	28.1	35.1	28.3	37.3	33.4	36.7	30.2
Family Conflict	41.3	34.4	34.2	32.8	32.6	28.2	29.4	26.9	35.7	29.6	36.9	33.0	29.6	30.1	32.2	30.9	34.9	30.6	33.1	30.9
Family History of Antisocial Behavior	34.2	35.4	35.4	27.8	32.8	26.3	28.6	21.4	30.0	30.8	34.6	24.5	29.9	25.8	32.4	27.0	31.8	29.6	32.8	25.1
Parental Attitudes Favorable to ASB	30.9	35.4	23.3	20.2	40.9	48.3	32.0	28.7	45.2	54.0	36.5	32.1	45.3	49.8	36.8	33.2	40.4	46.8	32.1	28.4
Parental Attitudes Favorable to Drug Use	10.3	9.4	5.0	3.9	17.6	19.3	11.1	9.3	27.3	27.3	21.2	14.7	23.8	23.1	20.1	13.2	19.5	19.7	14.4	10.2
<b>School Domain</b>																				
Academic Failure	41.3	38.0	37.1	29.2	40.5	37.8	38.0	28.6	40.8	38.9	38.6	30.8	36.2	33.8	44.8	33.5	39.7	37.2	39.6	30.5
Low Commitment to School	44.0	38.2	33.2	34.1	42.8	35.3	39.6	39.5	41.3	32.3	42.1	35.9	38.1	34.7	42.1	38.2	41.6	35.1	39.3	36.9
<b>Peer-Individual Domain</b>																				
Rebelliousness	25.5	24.7	16.3	16.1	32.4	26.7	26.1	21.7	35.9	32.0	36.5	29.0	41.0	32.2	28.2	28.5	33.5	29.0	26.9	23.7
Early Initiation of ASB	23.8	23.9	21.3	18.5	31.8	28.0	31.7	25.0	39.7	33.9	35.5	28.8	34.5	35.8	34.1	28.4	32.2	30.2	30.7	25.0
Early Initiation of Drug Use	18.0	14.6	16.9	11.1	25.2	22.0	18.1	15.4	23.1	20.6	24.3	16.3	20.2	22.4	24.1	16.9	21.6	19.8	20.9	14.9
Attitudes Favorable to ASB	32.1	25.5	30.6	27.7	32.7	23.9	26.9	25.9	41.8	34.2	34.8	31.0	44.2	31.0	37.4	32.3	37.5	28.6	32.4	29.1
Attitudes Favorable to Drug Use	12.8	7.8	10.2	8.4	23.8	19.3	22.0	18.0	31.3	25.7	30.9	24.4	25.9	22.4	30.1	22.0	23.2	18.7	23.3	18.1
Perceived Risk of Drug Use	36.1	40.5	34.9	31.0	26.2	23.2	27.8	24.6	39.0	37.0	37.3	31.2	31.9	30.7	34.0	28.9	33.3	32.8	33.5	28.9
Interaction with Antisocial Peers	35.4	38.8	28.9	22.9	32.8	27.7	25.5	20.5	33.0	30.1	26.7	21.6	30.7	30.3	21.6	20.4	33.0	31.7	25.7	21.3
Friend's Use of Drugs	15.1	11.9	10.4	8.5	31.5	25.5	24.4	18.0	27.2	24.9	26.7	18.3	20.0	24.5	20.2	15.1	23.4	21.7	20.4	14.9
Rewards for ASB	17.8	24.3	21.2	18.4	30.5	26.4	30.3	26.5	35.6	30.8	32.9	27.3	30.2	30.3	31.4	29.6	28.2	28.0	29.0	25.5
Depressive Symptoms	41.5	37.3	34.1	29.9	42.3	38.5	37.5	34.1	37.8	36.7	38.7	39.2	33.7	32.7	33.6	35.0	39.0	36.4	36.0	34.5
Gang Involvement	3.3	2.4	3.4	2.5	7.0	3.9	3.3	3.6	6.2	3.7	2.9	3.5	6.0	6.1	2.3	3.0	5.6	4.0	3.0	3.1

# Data Tables

**Table 9. Percentage of Students Reporting Protection**

Protective Factor	Grade 6				Grade 8				Grade 10				Grade 12				All Grades			
	LSAA 2009	LSAA 2011	LSAA 2013	State 2013																
<b>Community Domain</b>																				
Rewards for Prosocial Involvement	50.5	40.9	59.0	63.5	53.5	57.0	60.6	66.0	48.9	51.9	53.6	61.5	59.6	51.2	54.5	62.1	53.1	50.4	56.9	63.3
<b>Family Domain</b>																				
Family Attachment	63.1	56.9	67.9	70.4	58.8	63.9	66.4	69.1	65.3	65.3	64.3	71.3	70.3	70.4	70.3	71.1	64.5	64.1	67.3	70.5
Opportunities for Prosocial Involvement	63.8	57.0	65.2	68.4	65.3	67.6	70.1	73.0	65.9	57.1	61.4	67.6	70.6	66.3	63.3	66.8	66.4	62.0	65.0	69.0
Rewards for Prosocial Involvement	58.0	49.4	60.8	65.5	47.8	53.9	55.9	59.0	57.5	55.2	61.9	64.5	63.1	63.7	60.7	62.5	56.8	55.5	59.9	62.9
<b>School Domain</b>																				
Opportunities for Prosocial Involvement	43.5	55.1	53.8	59.9	64.2	72.3	72.6	70.7	67.6	71.4	70.4	74.4	70.4	72.0	71.4	75.8	61.1	67.6	67.1	70.0
Rewards for Prosocial Involvement	59.7	65.7	70.5	68.7	51.3	58.2	58.9	58.3	63.5	67.4	67.1	68.8	63.0	59.1	52.9	54.8	59.2	62.7	62.3	62.8
<b>Peer-Individual Domain</b>																				
Religiosity	43.8	43.9	45.5	55.7	57.0	64.2	58.3	69.7	56.1	56.7	57.5	66.4	64.0	60.4	51.7	67.2	55.2	56.5	53.2	64.8
Belief in the Moral Order	64.2	61.3	63.3	70.4	64.4	66.5	67.0	73.7	56.1	54.3	54.0	62.2	55.8	59.5	55.5	61.5	60.3	60.2	59.6	67.0
Interaction with Prosocial Peers	53.9	58.2	52.6	54.0	60.8	68.7	58.7	66.1	57.4	66.9	60.4	70.0	62.3	61.5	60.9	69.3	58.6	63.9	58.1	64.7
Prosocial Involvement	54.7	61.1	48.8	55.7	57.7	64.4	55.1	59.0	51.3	55.9	53.5	61.2	56.6	61.0	52.3	62.1	55.1	60.6	52.4	59.4
Rewards for Prosocial Involvement	49.0	55.3	62.2	61.6	57.1	62.6	64.5	64.4	59.8	70.9	70.5	74.6	71.9	76.5	70.9	74.9	59.2	66.5	67.2	68.9

# Data Tables

**Table 10. Drug Free Communities Report (2013 data)**

Outcome	Definition	Substance	LSAA 2013													
			Grade 6		Grade 8		Grade 10		Grade 12		All Grades**		Male †		Female †	
			Percent	Sample	Percent	Sample	Percent	Sample	Percent	Sample	Percent	Sample	Percent	Sample	Percent	Sample
<b>Perception of Risk*</b> (People are at Moderate or Great Risk of harming themselves if they...)	have five or more drinks of an alcoholic beverage once or twice a week	Alcohol	80.5	1,139	84.0	1,059	84.3	772	85.3	708	83.5	3,678	81.9	1,823	85.1	1,844
	smoke 1 or more packs of cigarettes per day	Cigarettes	90.5	1,139	90.8	1,065	93.6	783	94.3	725	92.3	3,712	92.1	1,839	92.6	1,861
	smoke marijuana once or twice a week	Marijuana	81.8	1,140	77.6	1,060	69.3	767	64.5	702	73.4	3,669	70.4	1,816	76.4	1,842
	use prescription drugs not prescribed to them	Prescriptions	88.6	1,135	91.7	1,059	93.2	764	93.5	698	91.7	3,656	90.9	1,813	92.6	1,831
<b>Perception of Parental Disapproval*</b> (Parents feel it would be Wrong or Very Wrong to...)	have one or two drinks of an alcoholic beverage nearly every day	Alcohol	98.7	1,135	98.1	1,062	97.9	782	96.2	723	97.7	3,702	97.1	1,833	98.4	1,858
	smoke tobacco	Tobacco	99.7	1,137	99.0	1,061	98.0	785	96.0	723	98.2	3,706	98.1	1,834	98.2	1,861
	smoke marijuana	Marijuana	98.9	1,134	97.4	1,060	95.7	784	92.9	722	96.2	3,700	95.8	1,833	96.7	1,856
	use prescription drugs not prescribed to you	Prescriptions	99.2	1,131	98.4	1,059	98.5	782	98.2	723	98.6	3,695	98.9	1,828	98.2	1,856
<b>Perception of Peer Disapproval*</b> (Friends feel it would be Wrong or Very Wrong to...)	have one or two drinks of an alcoholic beverage nearly every day	Alcohol	96.7	1,135	89.3	1,055	83.3	781	74.9	722	86.0	3,693	85.4	1,824	86.6	1,858
	smoke tobacco	Tobacco	97.5	1,135	93.4	1,054	88.4	781	78.8	722	89.5	3,692	88.3	1,823	90.7	1,858
	smoke marijuana	Marijuana	97.1	1,135	84.7	1,051	73.3	779	67.1	721	80.5	3,686	79.3	1,820	81.7	1,856
	use prescription drugs not prescribed to you	Prescriptions	97.9	1,134	94.3	1,052	92.9	778	87.0	722	93.0	3,686	92.4	1,822	93.7	1,854
<b>Past 30-Day Use*</b>	at least one use in the past 30 days	Alcohol	1.9	1,135	7.2	1,067	13.3	779	21.1	716	10.9	3,697	10.0	1,830	11.8	1,855
		Tobacco	0.9	1,063	3.2	960	4.9	714	8.7	665	4.5	3,402	5.0	1,656	3.9	1,737
		Marijuana	1.1	1,142	5.3	1,068	12.1	778	14.4	717	8.2	3,705	9.2	1,838	7.2	1,854
		Prescriptions	1.1	1,144	2.3	1,077	3.3	786	6.9	726	3.4	3,733	3.2	1,852	3.7	1,869

\* For Past 30-Day Use, Perception of Risk, and Perception of Parental/Peer Disapproval, the "Sample" column represents the sample size - the number of people who answered the question and whose responses were used to determine the percentage. The "Percent" column represents the percentage of youth in the sample answering the question as specified in the definition.

\*\* "All Grades" represents responses from students in all grades surveyed. The "All Grades" sample may contain additional data from grades that did not make the sample cutoff, and so may exceed the sum of the individual grade columns displayed. (In order to report individual grades/genders accurately, the grade or gender must have a minimum of twenty students reporting data. "All Grades" data not meeting the minimum number of respondents are displayed as "n/a.")

† The male and female values allow a gender comparison for youth who completed the survey. However, unless the percentage of students who participated from each grade is similar, the gender results are not necessarily representative of males and females in the community. In order to preserve confidentiality, male or female values may be omitted if the total number surveyed for that gender is under 20.

# Data Tables

**Table 11. Additional Data for Prevention Planning**

		Grade 6				Grade 8				Grade 10				Grade 12				All Grades			
		LSAA 2009	LSAA 2011	LSAA 2013	State 2013																
<b>Safety</b>																					
During the past 30 days, on how many days did you not go to school because you felt you would be unsafe at school or on your way to school?	One Or More Days	8.9	8.7	12.1	9.1	12.6	6.7	10.6	9.1	8.2	5.7	11.3	7.4	4.7	3.9	12.0	7.4	8.5	6.2	11.5	8.3
During the past 12 months, how often have you been picked on or bullied by a student ON SCHOOL PROPERTY?	More Than Once	29.7	21.4	30.6	28.5	21.5	17.1	30.2	26.4	12.3	12.8	18.1	17.3	8.6	6.7	9.8	11.8	17.9	14.2	22.1	21.2
<b>Discipline</b>																					
My teachers maintain good discipline in the classroom.	Strongly Agree or Agree	91.5	90.7	92.9	92.9	82.8	88.8	88.4	89.4	83.8	86.0	86.6	89.1	88.1	85.5	89.5	90.3	86.7	87.6	89.4	90.5
The principle and assistant principal maintain good discipline at my school.	Strongly Agree or Agree	86.9	87.8	89.7	89.3	83.1	85.5	85.2	87.1	80.2	85.1	87.5	87.8	84.0	84.5	87.7	87.1	83.6	85.7	87.5	87.9
<b>Perceived vs. Actual ATOD Use</b>																					
Smoke Cigarettes every day	Perceived Use	2.2	2.2	3.0	3.1	17.0	17.8	18.7	15.9	28.5	30.1	29.2	23.9	27.4	30.8	27.4	23.8	18.2	20.1	19.6	16.5
	Actual Use	0.3	0.1	0.4	0.2	1.3	1.1	1.0	0.4	4.1	3.7	1.5	1.6	2.4	6.1	4.6	2.5	1.9	2.7	1.9	1.1
Drank Alcohol in past 30 days	Perceived Use	4.6	2.9	4.5	3.9	26.1	25.2	25.2	20.1	43.4	42.0	41.5	33.8	46.2	46.5	46.9	37.7	29.2	29.0	29.7	23.6
	Actual Use	2.1	0.8	1.9	1.0	10.1	7.9	7.2	4.2	18.1	12.8	13.3	9.4	20.3	20.2	21.1	14.0	12.4	10.1	10.9	7.0
Used Marijuana in past 30 days	Perceived Use	1.9	1.1	3.0	2.5	18.2	19.2	24.3	19.4	31.8	34.5	39.0	31.7	33.2	36.0	37.8	33.4	20.6	22.7	26.1	21.5
	Actual Use	0.7	0.3	1.1	0.6	5.0	4.1	5.3	4.2	10.3	10.0	12.1	9.1	7.7	12.6	14.4	9.9	5.8	6.6	8.2	5.8

## Substance Use and Perceived Parental Acceptability

**Table 12. Substance Use in Relation to Perceived Parental Acceptability (State 2013)**

How wrong do your parents feel it would be for YOU to:	Student has used:	
	Alcohol At Least Once in Lifetime	Alcohol At Least Once in Past 30 Days
drink beer, wine, or hard liquor regularly?		
Very Wrong	15.4	4.5
Wrong	64.6	29.1
A Little Bit Wrong	81.4	44.1
Not Wrong At All	65.8	44.2
smoke marijuana?	Marijuana At Least Once in Lifetime	Marijuana At Least Once in Past 30 Days
Very Wrong	9.7	3.7
Wrong	49.9	25.9
A Little Bit Wrong	72.2	46.5
Not Wrong At All	74.3	53.6
smoke cigarettes?	Cigarettes At Least Once in Lifetime	Cigarettes At Least Once in Past 30 Days
Very Wrong	10.0	1.9
Wrong	43.9	15.1
A Little Bit Wrong	73.3	41.4
Not Wrong At All	47.4	34.0
use prescription drugs not prescribed to you?	Prescription Drugs At Least Once in Lifetime	Prescription Drugs At Least Once in Past 30 Days
Very Wrong	5.3	1.8
Wrong	25.4	11.3
A Little Bit Wrong	43.7	19.6
Not Wrong At All	33.8	14.0

## Even a Small Amount of Perceived Parental Acceptability Can Lead to Substance Use

When parents have favorable attitudes toward drugs, they influence the attitudes and behavior of their children. For example, parental approval of moderate drinking, even under parental supervision, substantially increases the risk of the young person using alcohol. Further, in families where parents involve children in their own drug or alcohol behavior, for example, asking the child to light the parent's cigarette or to get the parent a beer, there is an increased likelihood that their children will become drug users in adolescence.

In the Utah PNA Survey, students were asked how wrong their parents felt it was to use alcohol, marijuana, cigarettes, or prescription drugs not prescribed to them. The tables above display lifetime and past 30 days use rates in relation to parents'

acceptance of alcohol, marijuana, cigarette, or prescription drug use.

As Table 12 shows, relatively few students (15.4% lifetime, 4.5% 30-day) use alcohol when their parents think it is "Very Wrong" to use it. In contrast, when a student believes that their parents agree with use somewhat (i.e. the parent only believes that it is "Wrong," not "Very Wrong"), alcohol use increases to 64.6% for lifetime use and 29.1% for 30-day use. Similar findings can be observed regarding marijuana, cigarette and prescription drug use.

Table 12 illustrates how even a small amount of perceived parental acceptability can lead to substance use. These results make a strong argument for the importance of parents having strong and clear standards and rules when it comes to ATOD use.

## Appendix: Changes in the 2013 PNA

As new issues come to the forefront and new prevention modalities are implemented, the SHARP PNA survey evolves to reflect these concerns.

### Weighting procedures for 2013

During the analysis of SHARP survey data, Bach Harrison analysts have applied weights to the data to make the results more accurately reflect the total population of Utah students in grades 6, 8, 10, and 12.

Beginning in 2011, the State of Utah requested that Bach Harrison change the weighting procedure to account for the probability of a school participating in the survey and the probability of inclusion of students in each grade and gender category in each school. This differed from the weighting procedure used with past SHARP surveys (2009 and earlier) that was based upon school district enrollment in grades 6, 8, 10, and 12.

Comparison of the weighting procedures (2011/2013 vs. 2009 and earlier) produced comparable data.

### Changes to ATOD Questions

For the 2013 SHARP PNA, lifetime use is calculated from questions asking about age of first use; previous years are based off of the number of occasions used. 2013 lifetime use counts were obtained by generating a count of students answering any response other than *Never* to the question "How old were you when you first..." (used marijuana, used inhalants, etc.). In previous surveys, these data were obtained by counting the number of students having indicated one or more occasions of use of the substance in their lifetime. Significant analysis was conducted prior to the switch and Bach Harrison found that the two methods gathered comparable data; however, report readers should keep this change in mind as they compare 2013 data for lifetime use to previous years' data. The switch allowed removal of redundant questions, freeing up survey space and reducing survey completion time without sacrificing lifetime use data.

Lifetime use of alcohol and tobacco are exceptions to this change. Since several agencies track alcohol and tobacco use, lifetime use of these substances is calculated using separate questions (identical to previous years) to ensure that the results continue to be directly comparable from one administration to the next.

The change in calculating lifetime use resulted in a slight change to the way drug treatment needs was calculated. As with previous surveys, the "Needs Drug Treatment" continues to require that students answer *YES* to at least 3 drug treatment questions, but now requires any lifetime drug use, rather than drug use on 10 or more occasions.

Prescription Drugs is a new measure calculated by combining the responses of Prescription Stimulant, Prescription Sedative, Prescription Tranquilizer, and Prescription Narcotic Prescription Drugs.

Drug Free Communities measures were amended to conform to updated reporting requirements. New questions pertain to perceived risks, parental disapproval, and peer disapproval of alcohol, tobacco, marijuana, and prescription drug use.

In January 2014 SHARP reports were revised to correct lifetime use rates for substances other than alcohol and tobacco. While the corrections mainly affect the results for grades 10, 12, and all grades combined, some very small changes may be seen in the rates for grades 6 and 8.

In addition, lifetime and 30-day E-Cigarette data (available for 2011 and 2013) have been added in the January 2014 revised reports.

### New Health-Related Questions

Extra tobacco and health department questions were added in 2013. These include questions about:

- 1) student seatbelt use
- 2) participation in muscle strengthening activities
- 3) type of alcohol students used
- 4) days of school missed due to asthma
- 5) whether students had an asthma plan
- 6) students who had ridden in a car with a driver who was on a cell phone
- 7) exposure to tobacco advertisements

### Changes to Alcohol Questions

In 2011, all questions related to the sources of obtaining alcohol were removed. These questions were restored in 2013. The places of alcohol use were the same as prior years. A question was also added to regarding parental permissiveness of drinking alcohol.

## Appendix: Changes in the 2013 PNA (cont'd)

### Other Survey Removals and Changes

The number of occasions of lifetime substance use questions were removed (as noted above), with the exception of lifetime alcohol and tobacco use.

Other questions removed included questions about:

- 1) student consumption of sweetened drinks
- 2) hours of TV watched
- 3) use of flavored chewing tobacco
- 4) where students bought their last pack of cigarettes
- 5) if students were living with someone who used chewing tobacco.
- 6) student entitlement (“I deserve more things in life,” “Things should go my way.”)
- 7) the police catching someone drinking and driving in their neighborhood
- 8) *Intention to Use* risk factor scale was removed.

Finally, modifications were made in the order and inclusion of some questions on one or both forms in order to keep the form at a length suitable for administration during a 50 minute class period.

# Contacts for Prevention

## National Contacts

National Institute on Alcohol  
Abuse and Alcoholism  
<http://www.niaaa.nih.gov>

National Clearinghouse for  
Alcohol and Drug Information  
<http://store.samhsa.gov/>

The National Institute on Drug Abuse (NIDA)  
Drugs of Abuse Information Clearinghouse  
<http://www.nida.nih.gov/DrugPages.html>

Center for Substance Abuse Prevention  
<http://www.samhsa.gov/prevention/>

Monitoring the Future  
<http://monitoringthefuture.org>

National Survey on Drug Use and Health  
<http://www.oas.samhsa.gov/nsduh.htm>

## State Contacts

Utah Division of Substance Abuse  
and Mental Health  
120 North 200 West, #209  
Salt Lake City, Utah 84103  
<http://dsamh.utah.gov/>

Craig L. PoVey, Program Administrator  
801-538-4354  
Email: [clpovey@utah.gov](mailto:clpovey@utah.gov)

Ben Reaves, Program Manager  
801-538-3946  
Email: [breaves@Utah.gov](mailto:breaves@Utah.gov)

Brenda Ahlemann, Research Consultant  
801-538-9868  
Email: [bahlemann@utah.gov](mailto:bahlemann@utah.gov)

Susannah Burt, Program Manager  
801-538-4388  
Email: [sburt@utah.gov](mailto:sburt@utah.gov)

Utah State Office of Education  
Verne Larsen  
Coordinator, At Risk Services  
250 East 500 South  
Salt Lake City, Utah 84111  
801-538-7583  
Email: [larsen.verne@schools.utah.gov](mailto:larsen.verne@schools.utah.gov)

Utah Department of Health  
Janae Duncan  
Tobacco Prevention and Control Program  
801 538-9273  
288 North 1460 West  
Salt Lake City, UT 84116  
Email: [janaeduncan@utah.gov](mailto:janaeduncan@utah.gov)

## Regional Contacts

Bear River Planning District  
Billy Reamer  
Bear River Health Department  
655 East 1300 North  
Logan, UT 84341  
435-792-6529  
E-Mail: [breamer@brhd.org](mailto:breamer@brhd.org)

Central Planning District  
Sharon Lopez  
Central Utah Counseling Center  
255 South Main  
Richfield, UT 84701  
435-896-8236  
E-Mail: [sharonl@cucc.us](mailto:sharonl@cucc.us)

Davis Planning District  
Debi Todd  
Davis Behavioral Health  
2250 N. 1700 W.  
Layton, UT 84041  
801-447-8459  
E-Mail: [debit@dbhutah.org](mailto:debit@dbhutah.org)

## Contacts for Prevention

### Regional Contacts (Cont.)

#### Four Corners Planning District

Meranda Saccomano  
Four Corners Behavioral Health  
575 West 100 North  
Price, UT 84501  
435-637-2358, ext 220  
E-Mail: [msaccomano@fourcorners.ws](mailto:msaccomano@fourcorners.ws)

#### Northeastern Planning District

Robin Hatch (Vice Chair)  
Northeastern Counseling Center  
285 W. 800 S.  
Roosevelt, UT 84066  
435-725-6334  
E-Mail: [robinh@nccutah.org](mailto:robinh@nccutah.org)

#### Salt Lake Planning District

Jeff Smart & Kitt Curtis  
Salt Lake County Government Center  
2001 S. State Suite S-2300  
Salt Lake City, UT 84190  
801-468-2042 (Jeff) / 801-468-2031 (Kitt)  
E-Mail: [jsmart@SLCo.org](mailto:jsmart@SLCo.org)  
E-Mail: [kcurtis@slco.org](mailto:kcurtis@slco.org)

#### San Juan Planning District

Leslie Wojcik  
San Juan Counseling  
356 S. Main  
Blanding, UT 84511  
435-678-3262  
E-Mail: [lwojcik@sanjuancc.org](mailto:lwojcik@sanjuancc.org)

#### Southwest Planning District

Allen Sain  
Southwest Center  
245 East 680 South  
Cedar City, UT 84720  
435-867-7622  
E-Mail: [asain@sbhcutah.org](mailto:asain@sbhcutah.org)

#### Summit Planning District

Pamella Bello  
Valley Mental Health  
1753 Sidewinder Drive  
Park City, UT 84060  
435-649-8347  
E-Mail: [pamellab@vmh.com](mailto:pamellab@vmh.com)

#### Tooele Planning District

Julie Spindler  
Valley Mental Health  
100 South 1000 West  
Tooele, UT 84074  
435-843-3538  
E-Mail: [julies@vmh.com](mailto:julies@vmh.com)

#### Utah County Planning District

Pat Bird  
Utah County Dept of Drug & Alcohol Prevention  
& Treatment  
151 South University Avenue Suite 3200  
Provo, UT 84601  
801-851-7126  
E-Mail: [patbi@utahcounty.gov](mailto:patbi@utahcounty.gov)

#### Wasatch Planning District

Colleen Oshier  
Wasatch Mental Health  
55 South 500 East  
Heber, UT 84032  
435-654-3003  
E-Mail: [coshier@wasatch.org](mailto:coshier@wasatch.org)

#### Weber Planning District

Jennifer Hogge  
Weber Human Services  
237 26th Street  
Ogden, UT 84401  
801-625-3679  
E-Mail: [jenniferh@weberhs.org](mailto:jenniferh@weberhs.org)

#### This Report Was Prepared

for the State of Utah by  
by Bach Harrison LLC  
<http://www.bach-harrison.com>  
R. Steven Harrison, Ph.D.  
R. Paris Bach-Harrison, B.F.A.  
Taylor C. Bryant, B.A.  
Mary VanLeeuwen Johnstun, M.A.